

2

SYSTEMS OPTIMIZATION LABORATORY
DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD, CALIFORNIA 94305-4022

AD-A220 819

**A Monotone Complementarity Problem
in Hilbert Space**

by
Jen-Chih Yao

TECHNICAL REPORT SOL 90-7

April 1990

DTIC
ELECTE
APR 24 1990
S D
CO

Research and reproduction of this report were partially supported by the National Science Foundation grant DMS-8913089; Office of Naval Research grant N00014-89-J-1659, and the Department of Energy grant DE-FG03-87ER25028.

Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author and do NOT necessarily reflect the views of the above sponsors.

Reproduction in whole or in part is permitted for any purposes of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

90 04 20 053

A Monotone Complementarity Problem in Hilbert Space

Jen-Chih Yao

April 1990

Department of Operations Research

Stanford University

Stanford, CA 94305-4022

Abstract

An existence theorem for a complementarity problem involving a weakly coercive monotone mapping over an arbitrary closed convex cone in a real Hilbert space is established.

1. Introduction

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. Let K be a nonempty subset of H and f be a mapping from K into H . f is said to be *weakly coercive* if

$$\langle x, f(x) \rangle \rightarrow \infty \quad \text{as } \|x\| \rightarrow \infty \text{ and } x \in K.$$

f is said to be *monotone* if

$$\langle x - y, f(x) - f(y) \rangle \geq 0 \quad \text{for all } x, y \in K.$$

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Availability or Special
A-1	

f is said to be *strictly monotone* if the above inequality is strict whenever x and y are distinct.

f is said to be *strongly monotone* if there exists a positive number c such that

$$\langle x - y, f(x) - f(y) \rangle \geq c\|x - y\|^2 \quad \text{for all } x, y \in K.$$

A subset K of a real Hilbert space H is said to be a *cone* if $\lambda x \in K$ for all $x \in K$ and all $\lambda \geq 0$. Let K be a closed convex cone in H and dual cone K^* , that is,

$$K^* = \{u \in H \mid \langle u, x \rangle \geq 0, \forall x \in K\}.$$

The *complementarity problem* (CP) is to find $x \in K$ such that

$$f(x) \in K^* \quad \text{and} \quad \langle x, f(x) \rangle = 0. \quad (1)$$

Problem (1) was formulated by Karamardian [7] and has been extensively studied in the literature. See, e.g., [2, 4, 5, 6, 7, 8] and the references therein. The purpose of this paper is to prove an existence theorem for a complementarity problem involving a weakly coercive monotone mapping over an arbitrary closed convex cone in a real Hilbert space. The main result extends some existing results; the method of the proof of this main result is to consider the family of finite-dimensional subspaces by using the known results for finite-dimensional spaces and to show that a certain net of solutions from such subspaces converges to a solution to CP.

2. The Main Result

Now we prove the main result.

Theorem 2.1. *Let K be a closed convex cone in the real Hilbert space H . Let f be a weakly coercive monotone mapping from K into H which is continuous on $K \cap U$ for any*

finite-dimensional subspace U of H . Then there exists $x \in K$ such that

$$f(x) \in K^* \text{ and } \langle x, f(x) \rangle = 0.$$

Proof. Let U be any finite-dimensional subspace of H with $K \cap U \neq \emptyset$ and let P_U be the orthogonal projection of H onto U . Let $f_U = P_U f$ be the composition of P_U and f . Since the adjoint P_U^* of P_U is itself, we have

$$\lim_{\|u\| \rightarrow \infty, u \in K \cap U} \langle u, f_U(u) \rangle = \infty.$$

Therefore f_U is weakly coercive on $K \cap U$. By [1, Corollary 2.3], there exists $x_U \in K \cap U$ such that

$$\langle u - x_U, f(x_U) \rangle \geq 0 \quad \text{for all } u \in K \cap U. \quad (2)$$

Then by [7, Lemma 3.1], we have

$$f_U(x_U) \in (K \cap U)^* \text{ and } \langle x_U, f(x_U) \rangle = 0. \quad (3)$$

Let Λ be the family of all finite-dimensional subspaces U of H with $K \cap U \neq \emptyset$ and $K_U = \{x_U \mid U \subset V \in \Lambda\}$. Since f is weakly coercive, it follows from (3) that there exists a constant $r > 0$ so that $K_U \subset \bar{B}_r$ for all $U \in \Lambda$ where \bar{B}_r is the closure of the ball with center at 0 and radius r . For $U \in \Lambda$, let \bar{K}_U^w be the weak closure of K_U . Then the family $\{\bar{K}_U^w \mid U \in \Lambda\}$ has the finite intersection property. Indeed, for $U, V \in \Lambda$, let $W \in \Lambda$ be such that $U \cup V \subset W$. Then $K_U \cap K_V \supset K_W \neq \emptyset$. Since \bar{B}_r is weakly compact and $\bar{K}_U^w \subset \bar{B}_r$ for all $U \in \Lambda$, it follows that $\bigcap_{U \in \Lambda} \bar{K}_U^w \neq \emptyset$.

Let $x \in \bigcap_{U \in \Lambda} \bar{K}_U^w$. Suppose $u \in K$ is arbitrary and let $U \in \Lambda$ contain u . Since K_U is bounded and $x \in \bar{K}_U^w$, there exists a sequence $\{x_n\} \subset K_U$ which converges to x weakly. Since f is monotone, by (2) we have

$$\langle u - x_n, f(u) \rangle \geq 0 \quad \text{for all } n.$$

Since $\langle u - \cdot, f(u) \rangle$ is weakly continuous, we have

$$\langle u - x, f(u) \rangle \geq 0 \quad \text{for all } u \in K.$$

For any $u \in K$ and any $0 < t \leq 1$, let $u_t = tu + (1 - t)x$. By substituting u_t into (2), we have

$$\langle u_t - x, f(u_t) \rangle \geq 0 \quad \text{for all } 0 < t \leq 1. \quad (4)$$

Letting t approach 0 in (4), we get

$$\langle u - x, f(x) \rangle \geq 0 \quad \text{for all } u \in K. \quad (5)$$

By (5) and [7, Lemma 3.1], it follows that

$$f(x) \in K^* \text{ and } \langle x, f(x) \rangle = 0.$$

The next corollary follows from Theorem 2.1 directly.

Corollary 2.2. *Let K be a closed convex cone in the real Hilbert space H . Let f be a mapping from K into H which is continuous on $K \cap U$ for any finite-dimensional subspace U of H . Then there exists a unique $x \in K$ such that*

$$f(x) \in K^* \text{ and } \langle x, f(x) \rangle = 0$$

under each of the following conditions:

1. *f is strictly monotone and weakly coercive,*
2. *f is strongly monotone.*

We note that Corollary 2.2.2 extends a result of Nanda and Nanda [8, Theorem] where f is assumed to be strongly monotone and Lipschitzian.

References

1. M. Aganagić, "Variational inequalities and generalized complementarity problems," Technical Report SOL 78-11, System Optimization Laboratory, Department of Operations Research, Stanford University, Stanford, California, 1978. [See also "Contributions to Complementarity Theory," Ph.D. thesis, Department of Operations Research, Stanford University, Stanford, California, July 1978.]
2. M. S. Bazaraa, J. J. Goode and M. Z. Nashed, A nonlinear complementarity problem in mathematical programming in Banach space, *Proc. Amer. Math. Soc.* **35** (1972), 165-170.
3. K. Deimling, "Nonlinear Functional Analysis," Springer-Verlag, Berlin, 1985.
4. G. Isac and M. Théra, Complementarity problem and the existence of the post-critical equilibrium state of a thin elastic plate, *J. Optim. Theory Appl.* **58** (1988), 241-257.
5. G. Isac, Nonlinear complementarity problem and Galerkin method, *J. Math. Anal. Appl.* **108** (1985), 563-574.
6. S. Karamardian, Complementarity problems over cones with monotone and pseudomonotone maps, *J. Optim. Theory Appl.* **18** (1976), 445-454.
7. S. Karamardian, Generalized complementarity problem, *J. Optim. Theory Appl.* **8** (1971), 161-168.
8. S. Nanda and S. Nanda, A nonlinear complementarity problem in mathematical programming in Hilbert space, *Bull. Austral. Math. Soc.* **20** (1979), 233-236.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SOL 90-7	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Monotone Complementarity Problem in Hilbert Space		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Jen-Chih Yao		8. CONTRACT OR GRANT NUMBER(s) N00014-89-J-1659
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Operations Research - SOL Stanford University Stanford, CA 94305-4022		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1111MA
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research - Dept. of the Navy 800 N. Quincy Street Arlington, VA 22217		12. REPORT DATE April 1990
		13. NUMBER OF PAGES 5 pp.
		14. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Hilbert space; complementarity problem; monotone mapping; weakly coercive mapping.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An existence theorem for a complementarity problem involving a weakly coercive monotone mapping over an arbitrary closed convex cone in a real Hilbert space is established.		